



Gen. Math. Notes, Vol. 25, No. 1, November 2014, pp.8-18
ISSN 2219-7184; Copyright ©ICSRS Publication, 2014
www.i-csrs.org
Available free online at <http://www.geman.in>

Cesàro Summability of Double Sequences of Sets

Fatih Nuray¹, Uğur Ulusu² and Erdinç Dünder³

^{1,2,3}Department of Mathematics, Faculty of Science and Literature
Afyon Kocatepe University, Afyonkarahisar, Turkey

¹E-mail: fnuray@aku.edu.tr

²E-mail: ulusu@aku.edu.tr

³E-mail: edundar@aku.edu.tr

(Received: 1-6-14 / Accepted: 21-7-14)

Abstract

In this paper, we study the concepts of Wijsman Cesàro summability and Wijsman lacunary convergence of double sequences of sets and investigate the relationship between them.

Keywords: *Lacunary sequence, Cesàro summability, double sequence of sets, Wijsman convergence.*

1 Introduction

The concept of convergence of sequences of numbers has been extended by several authors to convergence of sequences of sets (see, [3, 4, 5, 11, 16, 17, 18]). Nuray and Rhoades [11] extended the notion of convergence of set sequences to statistical convergence and gave some basic theorems. Ulusu and Nuray [15] defined the Wijsman lacunary statistical convergence of sequence of sets and considered its relation with Wijsman statistical convergence, which was defined by Nuray and Rhoades. Ulusu and Nuray [16] introduced the concept of Wijsman strongly lacunary summability for set sequences and discussed its relation with Wijsman strongly Cesàro summability.

Hill [8] was the first who applied methods of functional analysis to double sequences. Also, Kull [9] applied methods of functional analysis of matrix maps of double sequences. A lot of usefull developments of double sequences in summability methods, the reader may refer to [1, 10, 14, 19].

In this paper, we study the concepts of Wijsman Cesàro summability and Wijsman lacunary convergence of double sequences of sets and investigate the relationship between them.

2 Definitions and Notations

Now, we recall the basic definitions and concepts (See [1, 2, 3, 4, 5, 11, 12, 14, 16, 17, 18]).

For any point $x \in X$ and any non-empty subset A of X , we define the distance from x to A by

$$d(x, A) = \inf_{a \in A} \rho(x, a).$$

Throughout the paper, we let (X, ρ) be a metric space and A, A_k be any non-empty closed subsets of X .

We say that the sequence $\{A_k\}$ is Wijsman convergent to A if

$$\lim_{k \rightarrow \infty} d(x, A_k) = d(x, A)$$

for each $x \in X$. In this case we write $W - \lim A_k = A$.

The sequence $\{A_k\}$ is said to be Wijsman Cesàro summable to A if $\{d(x, A_k)\}$ Cesàro summable to $\{d(x, A)\}$; that is, for each $x \in X$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n d(x, A_k) = d(x, A).$$

The sequence $\{A_k\}$ is said to be Wijsman strongly Cesàro summable to A if $\{d(x, A_k)\}$ strongly Cesàro summable to $\{d(x, A)\}$; that is, for each $x \in X$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n |d(x, A_k) - d(x, A)| = 0.$$

The sequence $\{A_k\}$ is said to be Wijsman strongly p -Cesàro summable to A if $\{d(x, A_k)\}$ strongly p -Cesàro summable to $\{d(x, A)\}$; that is, for each p positive real number and for each $x \in X$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n |d(x, A_k) - d(x, A)|^p = 0.$$

By a lacunary sequence we mean an increasing integer sequence $\theta = \{k_r\}$ such that $k_0 = 0$ and $h_r = k_r - k_{r-1} \rightarrow \infty$ as $r \rightarrow \infty$. Throughout this paper the intervals determined by θ will be denoted by $I_r = (k_{r-1}, k_r]$, and ratio $\frac{k_r}{k_{r-1}}$

will be abbreviated by q_r .

Let $\theta = \{k_r\}$ be a lacunary sequence. We say that the sequence $\{A_k\}$ is Wijsman lacunary convergent to A for each $x \in X$,

$$\lim_r \frac{1}{h_r} \sum_{k \in I_r} d(x, A_k) = d(x, A).$$

In this case we write $A_k \rightarrow A(WN_\theta)$.

Let $\theta = \{k_r\}$ be a lacunary sequence. We say that the sequence $\{A_k\}$ is Wijsman strongly lacunary convergent to A for each $x \in X$,

$$\lim_r \frac{1}{h_r} \sum_{k \in I_r} |d(x, A_k) - d(x, A)| = 0.$$

In this case we write $A_k \rightarrow A([WN_\theta])$.

A double sequence $x = (x_{kj})_{k,j \in \mathbb{N}}$ of real numbers is said to be convergent to $L \in \mathbb{R}$ in Pringsheim's sense if for any $\varepsilon > 0$, there exists $N_\varepsilon \in \mathbb{N}$ such that $|x_{kj} - L| < \varepsilon$ whenever $k, j > N_\varepsilon$. In this case we write

$$P - \lim_{k,j \rightarrow \infty} x_{kj} = L \quad \text{or} \quad \lim_{k,j \rightarrow \infty} x_{kj} = L.$$

A double sequence $x = (x_{kj})$ of real numbers is said to be bounded if there exists a positive real number M such that $|x_{kj}| < M$ for all $k, j \in \mathbb{N}$. That is

$$\|x\|_\infty = \sup_{k,j} |x_{kj}| < \infty.$$

The double sequence $\theta = \{(k_r, j_s)\}$ is called double lacunary sequence if there exist two increasing sequence of integers such that

$$k_0 = 0, \quad h_r = k_r - k_{r-1} \rightarrow \infty \quad \text{as} \quad r \rightarrow \infty$$

and

$$j_0 = 0, \quad \bar{h}_u = j_u - j_{u-1} \rightarrow \infty \quad \text{as} \quad u \rightarrow \infty.$$

We use the following notations in the sequel:

$$k_{ru} = k_r j_u, \quad h_{ru} = h_r \bar{h}_u, \quad I_{ru} = \{(k, j) : k_{r-1} < k \leq k_r \quad \text{and} \quad j_{u-1} < j \leq j_u\},$$

$$q_r = \frac{k_r}{k_{r-1}} \quad \text{and} \quad q_u = \frac{j_u}{j_{u-1}}.$$

Lemma 2.1 [7, Lemma 3.2] *If b_1, b_2, \dots, b_n are positive real numbers, and if a_1, a_2, \dots, a_n are real numbers satisfying*

$$\frac{|a_1 + a_2 + \dots + a_n|}{b_1 + b_2 + \dots + b_n} > \varepsilon > 0,$$

then $|a_i|/b_i > \varepsilon$ for some i , where $1 \leq i \leq n$.

3 Main Results

Throughout the paper, A, A_{kj} denote any non-empty closed subsets of X .

Definition 3.1 *The double sequence $\{A_{kj}\}$ is Wijsman convergent to A if*

$$P - \lim_{k,j \rightarrow \infty} d(x, A_{kj}) = d(x, A) \quad \text{or} \quad \lim_{k,j \rightarrow \infty} d(x, A_{kj}) = d(x, A)$$

for each $x \in X$. In this case we write $W_2 - \lim A_{kj} = A$.

Example 3.2 *Let $X = \mathbb{R}^2$ and $\{A_{kj}\}$ be the following double sequence:*

$$A_{kj} = \left\{ (x, y) \in \mathbb{R}^2 : x^2 + (y - 1)^2 = \frac{1}{kj} \right\}.$$

This double sequence of sets is Wijsman convergent to the set $A = \{(0, 1)\}$.

Definition 3.3 *The double sequence $\{A_{kj}\}$ is said to be Wijsman Cesàro summable to A if $\{d(x, A_{kj})\}$ Cesàro summable to $\{d(x, A)\}$; that is, for each $x \in X$,*

$$\lim_{m,n \rightarrow \infty} \frac{1}{mn} \sum_{k,j=1,1}^{m,n} d(x, A_{kj}) = d(x, A).$$

In this case we write $A_{kj} \xrightarrow{(W_2\sigma_1)} A$.

Definition 3.4 *The double sequence $\{A_{kj}\}$ is said to be Wijsman strongly Cesàro summable to A if $\{d(x, A_{kj})\}$ strongly Cesàro summable to $\{d(x, A)\}$; that is, for each $x \in X$,*

$$\lim_{m,n \rightarrow \infty} \frac{1}{mn} \sum_{k,j=1,1}^{m,n} |d(x, A_{kj}) - d(x, A)| = 0.$$

In this case we write $A_{kj} \xrightarrow{[W_2\sigma_1]} A$.

Example 3.5 *Let $X = \mathbb{R}^2$ and define the double sequence $\{A_{kj}\}$ by*

$$A_{kj} = \begin{cases} \{(x, y) \in \mathbb{R}^2 : (x - 1)^2 + (y - 1)^2 = k\} & , \quad j = 1, \text{ for all } k \\ \{(x, y) \in \mathbb{R}^2 : (x - 1)^2 + (y - 1)^2 = j\} & , \quad k = 1, \text{ for all } j \\ \{(0, 0)\} & , \quad \text{otherwise.} \end{cases}$$

Then $\{A_{kj}\}$ is Wijsman convergent to the set $A = \{(0, 0)\}$ but

$$\lim_{m,n \rightarrow \infty} \frac{1}{mn} \sum_{k,j=1,1}^{m,n} d(x, A_{kj})$$

does not tend to a finite limit. Hence, $\{A_{kj}\}$ is not Wijsman Cesàro summable. Also, $\{A_{kj}\}$ is not Wijsman strongly Cesàro summable.

Definition 3.6 The double sequence $\{A_{kj}\}$ is said to be Wijsman strongly p -Cesàro summable to A if $\{d(x, A_{kj})\}$ strongly p -Cesàro summable to $\{d(x, A)\}$; that is, for each p positive real number and for each $x \in X$,

$$\lim_{m,n \rightarrow \infty} \frac{1}{mn} \sum_{k,j=1,1}^{m,n} |d(x, A_{kj}) - d(x, A)|^p = 0.$$

In this case we write $A_{kj} \xrightarrow{[W_2\sigma_p]} A$.

Definition 3.7 Let $\theta = \{(k_r, j_s)\}$ be a double lacunary sequence. The double sequence $\{A_{kj}\}$ is Wijsman lacunary convergent to A if for each $x \in X$,

$$\lim_{r,u \rightarrow \infty} \frac{1}{h_r \bar{h}_u} \sum_{k=k_{r-1}+1}^{k_r} \sum_{j=j_{u-1}+1}^{j_u} d(x, A_{kj}) = d(x, A).$$

In this case we write $A_{kj} \xrightarrow{(W_2N_\theta)} A$.

Definition 3.8 Let $\theta = \{(k_r, j_s)\}$ be a double lacunary sequence. The double sequence $\{A_{kj}\}$ is Wijsman strongly lacunary convergent to A if for each $x \in X$,

$$\lim_{r,u \rightarrow \infty} \frac{1}{h_r \bar{h}_u} \sum_{k=k_{r-1}+1}^{k_r} \sum_{j=j_{u-1}+1}^{j_u} |d(x, A_{kj}) - d(x, A)| = 0.$$

In this case we write $A_{kj} \xrightarrow{[W_2N_\theta]} A$.

Definition 3.9 Let $\theta = \{(k_r, j_s)\}$ be a double lacunary sequence. The double sequence $\{A_{kj}\}$ is Wijsman strongly p -lacunary convergent to A if for each p positive real number and for each $x \in X$,

$$\lim_{r,u \rightarrow \infty} \frac{1}{h_r \bar{h}_u} \sum_{k=k_{r-1}+1}^{k_r} \sum_{j=j_{u-1}+1}^{j_u} |d(x, A_{kj}) - d(x, A)|^p = 0.$$

In this case we write $A_{kj} \xrightarrow{[W_2^p N_\theta]} A$.

Theorem 3.10 For any double lacunary sequence θ , if $\liminf_r q_r > 1$ and $\liminf_u q_u > 1$, then $[W_2\sigma_1] \subseteq [W_2N_\theta]$.

Proof: Assume that $\liminf_r q_r > 1$ and $\liminf_u q_u > 1$. Then there exist $\lambda, \mu > 0$ such that $q_r \geq 1 + \lambda$ and $q_u \geq 1 + \mu$ for all $r, u \geq 1$, which implies that

$$\frac{k_r j_u}{h_r \bar{h}_u} \leq \frac{(1 + \lambda)(1 + \mu)}{\lambda \mu}.$$

Let $A_{kj} \xrightarrow{[W_2\sigma_1]} A$. We can write

$$\begin{aligned} \frac{1}{h_r \bar{h}_u} \sum_{k,j \in I_{ru}} |d(x, A_{kj}) - d(x, A)| &= \frac{1}{h_r \bar{h}_u} \sum_{i,s=1,1}^{k_r, j_u} |d(x, A_{is}) - d(x, A)| \\ &\quad - \frac{1}{h_r \bar{h}_u} \sum_{i,s=1,1}^{k_{r-1}, j_{u-1}} |d(x, A_{is}) - d(x, A)| \\ &= \frac{k_r j_u}{h_r \bar{h}_u} \left(\frac{1}{k_r j_u} \sum_{i,s=1,1}^{k_r, j_u} |d(x, A_{is}) - d(x, A)| \right) \\ &\quad - \frac{k_{r-1} j_{u-1}}{h_r \bar{h}_u} \left(\frac{1}{k_{r-1} j_{u-1}} \sum_{i,s=1,1}^{k_{r-1}, j_{u-1}} |d(x, A_{is}) - d(x, A)| \right). \end{aligned}$$

Since $A_{kj} \xrightarrow{[W_2\sigma_1]} A$, the terms

$$\frac{1}{k_r j_u} \sum_{i,s=1,1}^{k_r, j_u} |d(x, A_{is}) - d(x, A)| \quad \text{and} \quad \frac{1}{k_{r-1} j_{u-1}} \sum_{i,s=1,1}^{k_{r-1}, j_{u-1}} |d(x, A_{is}) - d(x, A)|$$

both tend to 0, and it follows that

$$\frac{1}{h_r \bar{h}_u} \sum_{k,j \in I_{ru}} |d(x, A_{kj}) - d(x, A)| \rightarrow 0,$$

that is, $A_{kj} \xrightarrow{[W_2N_\theta]} A$. Hence, $[W_2\sigma_1] \subseteq [W_2N_\theta]$.

Theorem 3.11 *For any double lacunary sequence θ , if $\limsup_r q_r < \infty$ and $\limsup_u q_u < \infty$ then $[W_2N_\theta] \subseteq [W_2\sigma_1]$.*

Proof: Assume that $\limsup_r q_r < \infty$ and $\limsup_u q_u < \infty$, then there exists $M, N > 0$ such that $q_r < M$ and $q_u < N$, for all r, u . Let $\{A_{kj}\} \in [W_2N_\theta]$ and $\varepsilon > 0$. Then we can find $R, U > 0$ and $K > 0$ such that

$$\sup_{i \geq R, s \geq U} \tau_{is} < \varepsilon \quad \text{and} \quad \tau_{is} < K \quad \text{for all } i, s = 1, 2, \dots,$$

where

$$\tau_{ru} = \frac{1}{h_r \bar{h}_u} \sum_{I_{ru}} |d(x, A_{kj}) - d(x, A)|.$$

If t, v are any integers with $k_{r-1} < t \leq k_r$ and $j_{u-1} < v \leq j_u$, where $r > R$ and $u > U$, then we can write

$$\begin{aligned}
\frac{1}{tv} \sum_{i,s=1,1}^{t,v} |d(x, A_{is}) - d(x, A)| &\leq \frac{1}{k_{r-1}j_{u-1}} \sum_{i,s=1,1}^{k_r,j_u} |d(x, A_{is}) - d(x, A)| \\
&= \frac{1}{k_{r-1}j_{u-1}} \left(\sum_{I_{11}} |d(x, A_{is}) - d(x, A)| \right. \\
&\quad + \sum_{I_{12}} |d(x, A_{is}) - d(x, A)| \\
&\quad + \sum_{I_{21}} |d(x, A_{is}) - d(x, A)| \\
&\quad + \sum_{I_{22}} |d(x, A_{is}) - d(x, A)| \\
&\quad \left. + \cdots + \sum_{I_{ru}} |d(x, A_{is}) - d(x, A)| \right) \\
&\leq \frac{k_1 j_1}{k_{r-1} j_{u-1}} \cdot \tau_{11} + \frac{k_1 (j_2 - j_1)}{k_{r-1} j_{u-1}} \cdot \tau_{12} \\
&\quad + \frac{(k_2 - k_1) j_1}{k_{r-1} j_{u-1}} \cdot \tau_{21} \\
&\quad + \frac{(k_2 - k_1) (j_2 - j_1)}{k_{r-1} j_{u-1}} \cdot \tau_{22} \\
&\quad + \cdots + \frac{(k_R - k_{R-1}) (j_U - j_{U-1})}{k_{r-1} j_{u-1}} \tau_{RU} \\
&\quad + \cdots + \frac{(k_r - k_{r-1}) (j_u - j_{u-1})}{k_{r-1} j_{u-1}} \tau_{ru} \\
&\leq \left(\sup_{i,s \geq 1,1} \tau_{is} \right) \frac{k_R j_U}{k_{r-1} j_{u-1}} \\
&\quad + \left(\sup_{i \geq R, s \geq U} \tau_{is} \right) \frac{(k_r - k_R) (j_u - j_U)}{k_{r-1} j_{u-1}} \\
&\leq K \frac{k_R j_U}{k_{r-1} j_{u-1}} + \varepsilon M N.
\end{aligned}$$

Since $k_{r-1}, j_{u-1} \rightarrow \infty$ as $t, v \rightarrow \infty$, it follows that

$$\frac{1}{tv} \sum_{i,s=1,1}^{t,v} |d(x, A_{is}) - d(x, A)| \rightarrow 0$$

and consequently $\{A_{kj}\} \in [W_2\sigma_1]$. Hence, $[W_2N_\theta] \subseteq [W_2\sigma_1]$.

Theorem 3.12 *For any double lacunary sequence θ , if $1 < \liminf_r q_r \leq \limsup_r q_r < \infty$ and $1 < \liminf_u q_u \leq \limsup_u q_u < \infty$, then $[W_2N_\theta] = [W_2\sigma_1]$.*

Proof: This follows from Theorem 3.10 and Theorem 3.11.

Theorem 3.13 *For any double lacunary sequence θ , let $\{A_{kj}\} \in [W_2N_\theta] \cap [W_2\sigma_1]$. If $A_{kj} \xrightarrow{[W_2N_\theta]} A$ and $A_{kj} \xrightarrow{[W_2\sigma_1]} B$ then $A = B$.*

Proof: Let $A_{kj} \xrightarrow{[W_2\sigma_1]} A$, $A_{kj} \xrightarrow{[W_2N_\theta]} B$ and suppose that $A \neq B$. We can write

$$\begin{aligned} v_{ru} + \tau_{ru} &= \frac{1}{h_r \bar{h}_u} \sum_{k,j \in I_{ru}} |d(x, A_{kj}) - d(x, A)| + \frac{1}{h_r \bar{h}_u} \sum_{k,j \in I_{ru}} |d(x, A_{kj}) - d(x, B)| \\ &\geq \frac{1}{h_r \bar{h}_u} \sum_{k,j \in I_{ru}} |d(x, A) - d(x, B)| \\ &= |d(x, A) - d(x, B)|, \end{aligned}$$

where

$$v_{ru} = \frac{1}{h_r \bar{h}_u} \sum_{k,j \in I_{ru}} |d(x, A_{kj}) - d(x, A)| \text{ and } \tau_{ru} = \frac{1}{h_r \bar{h}_u} \sum_{k,j \in I_{ru}} |d(x, A_{kj}) - d(x, B)|.$$

Since $\{A_{kj}\} \in [W_2N_\theta]$, $\tau_{ru} \rightarrow 0$. Thus for sufficiently large r, u we must have

$$v_{ru} > \frac{1}{2} |d(x, A) - d(x, B)|.$$

Observe that

$$\begin{aligned} \frac{1}{k_r j_u} \sum_{i,s=1,1}^{k_r, j_u} |d(x, A_{is}) - d(x, A)| &\geq \frac{1}{k_r j_u} \sum_{I_{ru}} |d(x, A_{is}) - d(x, A)| \\ &= \frac{(k_r - k_{r-1})(j_u - j_{u-1})}{k_r j_u} \cdot v_{ru} \\ &= \left(1 - \frac{1}{q_r}\right) \left(1 - \frac{1}{q_u}\right) \cdot v_{ru} \\ &> \frac{1}{2} \left(1 - \frac{1}{q_r}\right) \left(1 - \frac{1}{q_u}\right) \cdot |d(x, A) - d(x, B)| \end{aligned}$$

for sufficiently large r, u . Since $\{A_{kj}\} \in [W_2\sigma_1]$, the left hand side of the inequality above convergent to 0, so we must have $q_r \rightarrow 1$ and $q_u \rightarrow 1$. But this implies, by proof of Theorem 3.11, that

$$[W_2N_\theta] \subset [W_2\sigma_1].$$

That is, we have

$$A_{kj} \xrightarrow{[W_2N_\theta]} B \Rightarrow A_{kj} \xrightarrow{[W_2\sigma_1]} B,$$

and therefore

$$\frac{1}{tv} \sum_{i,s=1,1}^{t,v} |d(x, A_{is}) - d(x, B)| \rightarrow 0.$$

Then, we have

$$\begin{aligned} \frac{1}{tv} \sum_{i,s=1,1}^{t,v} |d(x, A_{is}) - d(x, B)| + \frac{1}{tv} \sum_{i,s=1,1}^{t,v} |d(x, A_{is}) - d(x, A)| \\ \geq |d(x, A) - d(x, B)| > 0, \end{aligned}$$

which yields a contradiction to our assumption, since both terms on the left hand side tend to 0. That is, for each $x \in X$,

$$|d(x, A) - d(x, B)| = 0,$$

and therefore $A = B$.

Definition 3.14 *The double sequence $\theta' = \{(k'_r, j'_u)\}$ is called double lacunary refinement of the double lacunary sequence $\theta = \{(k_r, j_u)\}$ if $\{k_r\} \subseteq \{k'_r\}$ and $\{j_u\} \subseteq \{j'_u\}$.*

Theorem 3.15 *If θ' is a double lacunary refinement of double lacunary sequence θ and if $\{A_{kj}\} \notin [W_2N_\theta]$, then $\{A_{kj}\} \notin [W_2N_{\theta'}]$.*

Proof: Let $\{A_{kj}\} \notin [W_2N_\theta]$. Then, for any non-empty closed subset $A \subseteq X$ there exists $\varepsilon > 0$ and a subsequence (k_{r_n}) of (k_r) and (j_{u_n}) of (j_u) such that

$$\tau_{r_n u_n} = \frac{1}{h_{r_n} \bar{h}_{u_n}} \sum_{k,j=1,1}^{k_{r_n}, j_{u_n}} |d(x, A_{kj}) - d(x, A)| \geq \varepsilon.$$

Writing

$$I_{r_n u_n} = I'_{s+1,t+1} \cup I'_{s+1,t+2} \cup I'_{s+2,t+1} \cup I'_{s+2,t+2} \cup \dots \cup I'_{s+p,t+p}$$

where

$$k_{r_{n-1}} = k'_s < k'_{s+1} < \dots < k'_{s+p} = k_{r_n} \text{ and } j_{u_{n-1}} = j'_t < j'_{t+1} < \dots < j'_{t+p} = j_{u_n}.$$

Then we have

$$\tau_{r_n u_n} = \frac{\sum_{I'_{s+1,t+1}} |d(x, A_{k_j}) - d(x, A)| + \dots + \sum_{I'_{s+p,t+p}} |d(x, A_{k_j}) - d(x, A)|}{h'_{s+1} \bar{h}'_{t+1} + \dots + h'_{s+p} \bar{h}'_{t+p}}.$$

It follows from Lemma 2.1 that

$$\frac{1}{h'_{s+p} \bar{h}'_{t+p}} \sum_{I'_{s+p,t+p}} |d(x, A_{k_j}) - d(x, A)| \geq \varepsilon$$

for some j and consequently, $\{A_{k_j}\} \notin [W_2 N_{\theta'}]$.

References

- [1] B. Altay and F. Başar, Some new spaces of double sequences, *J. Math. Anal. Appl.*, 309(1) (2005), 70-90.
- [2] J.P. Aubin and H. Frankowska, *Set-Valued Analysis*, Birkhauser, Boston, (1990).
- [3] M. Baronti and P. Papini, Convergence of sequences of sets, In: *Methods of Functional Analysis in Approximation Theory*, ISNM 76, Birkhauser-Verlag, Basel, (1986), 133-155.
- [4] G. Beer, On convergence of closed sets in a metric space and distance functions, *Bull. Aust. Math. Soc.*, 31(1985), 421-432.
- [5] G. Beer, Wijsman convergence: A survey, *Set-Valued Var. Anal.*, 2(1994), 77-94.
- [6] J.S. Connor, The statistical and strong p-Cesaro convergence of sequences, *Analysis*, 8(1988), 46-63.
- [7] A.R. Freedman, J.J. Sember and M. Raphael, Some Cesaro type summability spaces, *Proc. London Math. Soc.*, 37(1978), 508-520.
- [8] J.D. Hill, On perfect summability of double sequences, *Bull. Amer. Math. Soc.*, 46(1940), 327-331.
- [9] I.G. Kull, Multiplication of summable double series, *Uch. zap. Tartusskogo un-ta*, 62(1958), 3-59 (in Russian).

- [10] B.V. Limayea and M. Zeltser, On the pringsheim convergence of double series, *Proc. Est. Acad. Sci.*, 58(2009), 108-121.
- [11] F. Nuray and B.E. Rhoades, Statistical convergence of sequences of sets, *Fasc. Math.*, 49(2012), 87-99.
- [12] A. Pringsheim, Zur theorie der zweifach unendlichen Zahlenfolgen, *Math. Ann.*, 53(1900), 289-321.
- [13] R.T. Rockafellar and R.J.B. Wets, *Variational Analysis*, Grundlehren der Mathematischen Wissenschaften 317, Springer-Verlag, (2009).
- [14] E. Savaş, On some double lacunary sequence spaces of fuzzy numbers, *Math. Comput. Appl.*, 15(3) (2010), 439-448.
- [15] U. Ulusu and F. Nuray, Lacunary statistical convergence of sequence of sets, *Prog. Appl. Math.*, 4(2) (2012), 99-109.
- [16] U. Ulusu and F. Nuray, On strongly lacunary summability of sequences of sets, *J. Appl. Math. Bioinform*, 3(3) (2013), 75-88.
- [17] R.A. Wijsman, Convergence of sequences of convex sets, cones and functions, *Bull. Amer. Math. Soc.*, 70(1964), 186-188.
- [18] R.A. Wijsman, Convergence of sequences of convex sets, cones and functions II, *Trans. Amer. Math. Soc.*, 123(1) (1966), 32-45.
- [19] M. Zeltser, M. Mursaleen and S.A. Mohiuddine, On almost conservative matrix methods for double sequence spaces, *Publ. Math. Debrecen*, 75(2009), 1-13.