

Wijsman \mathcal{I}_2 -Invariant Convergence of Double Sequences of Sets

Şükrü TORTOP and Erdiñç DÜNDAR

Department of Mathematics,
Afyon Kocatepe University,
03200 Afyonkarahisar, Turkey.

ICAA-2016
July 12–15, 2016

- 1 Abstract
- 2 Introduction
- 3 Definitions and Notations
- 4 Main Result
- 5 References

Abstract

In this paper, we study the concepts of Wijsman invariant convergence, Wijsman invariant statistical convergence, Wijsman \mathcal{I}_2 -invariant convergence ($\mathcal{I}_{W_2}^\sigma$), Wijsman \mathcal{I}_2^* -invariant convergence ($\mathcal{I}_{W_2}^{*\sigma}$), Wijsman p -strongly invariant convergence ($[W_2 V_\sigma]_p$) of double sequence of sets and investigate the relationships among Wijsman invariant convergence, $[W_2 V_\sigma]_p$, $\mathcal{I}_{W_2}^\sigma$ and $\mathcal{I}_{W_2}^{*\sigma}$. Also, we introduce the concepts of $\mathcal{I}_{W_2}^\sigma$ -Cauchy double sequence and $\mathcal{I}_{W_2}^{*\sigma}$ -Cauchy double sequence of sets.

Introduction

Throughout the paper \mathbb{N} denotes the set of all positive integers and \mathbb{R} the set of all real numbers. The concept of convergence of a sequence of real numbers has been extended to statistical convergence independently by Fast [6] and Schoenberg [30]. This concept was extended to the double sequences by Mursaleen and Edely [12]. Nuray and Ruckle [17] independently introduced the same with another name generalized statistical convergence. The idea of \mathcal{I} -convergence was introduced by Kostyrko, Šalát and Wilczyński [8] as a generalization of statistical convergence which is based on the structure of the ideal \mathcal{I} of subset of \mathbb{N} . Das et al. [4] introduced the concept of \mathcal{I} -convergence of double sequences in a metric space and studied some properties of this convergence. A lot of development have been made in this area after the works of [5, 9, 15].

Introduction

Nuray and Rhoades [16] extended the notion of convergence of set sequences to statistical convergence and gave some basic theorems. Ulusu and Nuray [35] defined the Wijsman lacunary statistical convergence of set sequences and considered its relation with Wijsman statistical convergence, which was defined by Nuray and Rhoades. Kişi and Nuray [7] introduced a new convergence notion, for sequences of sets, which is called Wijsman \mathcal{I} -convergence. Also, the concept of convergence of sequences has been extended to convergence, statistical convergence and ideal convergence of sequences of sets by several authors (see, [31, 32, 33, 34, 36, 37, 38, 39]).

Introduction

Several authors including Raimi [25], Schaefer [29], Mursaleen [14], Savaş [26], Pancaroğlu and Nuray [22], and others have studied invariant convergent sequences (see, [11, 19]). The concept of strongly σ -convergence was defined by Mursaleen [13]. Savaş and Nuray [28] introduced the concepts of σ -statistical convergence and lacunary σ -statistical convergence and gave some inclusion relations.

Introduction

Recently, the concept of strong σ -convergence was generalized by Savaş [26]. Nuray et al. [20] defined the concepts of σ -uniform density of subsets A of the set \mathbb{N} , \mathcal{I}_σ -convergence and investigated relationships between \mathcal{I}_σ -convergence and invariant convergence also \mathcal{I}_σ -convergence and $[V_\sigma]_p$ -convergence. Ulusu and Nuray [21] investigated lacunary \mathcal{I} -invariant convergence and lacunary \mathcal{I} -invariant Cauchy sequence of real numbers. Pancaroğlu et al. [24] studied Wijsman \mathcal{I} -invariant convergence of sequences of sets.

Definitions and Notations

A family of sets $\mathcal{I} \subseteq 2^{\mathbb{N}}$ is called an ideal if and only if

- (i) $\emptyset \in \mathcal{I}$,
- (ii) For each $A, B \in \mathcal{I}$ we have $A \cup B \in \mathcal{I}$,
- (iii) For each $A \in \mathcal{I}$ and each $B \subseteq A$ we have $B \in \mathcal{I}$.

An ideal is called nontrivial if $\mathbb{N} \notin \mathcal{I}$ and nontrivial ideal is called admissible if $\{n\} \in \mathcal{I}$ for each $n \in \mathbb{N}$.

A family of sets $\mathcal{F} \subseteq 2^{\mathbb{N}}$ is called a filter if and only if

- (i) $\emptyset \notin \mathcal{F}$,
- (ii) For each $A, B \in \mathcal{F}$ we have $A \cap B \in \mathcal{F}$,
- (iii) For each $A \in \mathcal{F}$ and each $B \supseteq A$ we have $B \in \mathcal{F}$.

Definitions and Notations

Proposition 1

([8]) \mathcal{I} is nontrivial ideal in \mathbb{N} if and only if
 $\mathcal{F}(\mathcal{I}) = \{M \subset \mathbb{N} : (\exists A \in \mathcal{I})(M = \mathbb{N} \setminus A)\}$ is a filter in \mathbb{N} .

A nontrivial ideal \mathcal{I}_2 of $\mathbb{N} \times \mathbb{N}$ is called strongly admissible ideal if $\{i\} \times \mathbb{N}$ and $\mathbb{N} \times \{i\}$ belong to \mathcal{I}_2 for each $i \in \mathbb{N}$.

Throughout the paper we take \mathcal{I}_2 as a strongly admissible ideal in $\mathbb{N} \times \mathbb{N}$. It is evident that a strongly admissible ideal is admissible also.

Definitions and Notations

$\mathcal{I}_2^0 = \{A \subset \mathbb{N} \times \mathbb{N} : (\exists m(A) \in \mathbb{N})(i, j \geq m(A) \Rightarrow (i, j) \notin A)\}$. Then \mathcal{I}_2^0 is a strongly admissible ideal and clearly an ideal \mathcal{I}_2 is strongly admissible if and only if $\mathcal{I}_2^0 \subset \mathcal{I}_2$.

Let (X, ρ) be a metric space. A sequence $x = (x_{mn})$ in X is said to be \mathcal{I}_2 -convergent to $L \in X$, if for any $\varepsilon > 0$,

$A(\varepsilon) = \{(m, n) \in \mathbb{N} \times \mathbb{N} : \rho(x_{mn}, L) \geq \varepsilon\} \in \mathcal{I}_2$. In this case, we say that x is \mathcal{I}_2 -convergent and we write $\mathcal{I}_2 - \lim_{m, n \rightarrow \infty} x_{mn} = L$.

Definitions and Notations

Let σ be a mapping of the set of positive integers into itself. A continuous linear functional ϕ on ℓ_∞ , the space of real bounded sequences, is said to be an invariant mean or a σ -mean if and only if

- 1 $\phi(x) \geq 0$, when the sequence $x = (x_n)$ has $x_n \geq 0$ for all n ,
- 2 $\phi(e) = 1$, where $e = (1, 1, 1, \dots)$, and
- 3 $\phi(x_{\sigma(n)}) = \phi(x_n)$ for all $x \in \ell_\infty$.

Definitions and Notations

The mappings σ are one-to-one and such that $\sigma^m(n) \neq n$ for all positive integers n and m , where $\sigma^m(n)$ denotes the m th iterate of the mapping σ at n . Thus ϕ extends the limit functional on c , the space of convergent sequences, in the sense that $\phi(x) = \lim x$ for all $x \in c$. In the case σ is translation mappings $\sigma(n) = n + 1$, the σ -mean is often called a Banach limit and V_σ , the set of bounded sequences all of whose invariant means are equal, is the set of almost convergent sequences [10]. It can be shown [27] that

$$V_\sigma = \left\{ x = (x_n) \in \ell_\infty : \lim_{m \rightarrow \infty} \frac{1}{m} \sum_{k=1}^m x_{\sigma^k(n)} = L \text{ uniformly in } n \right\}.$$

Definitions and Notations

A bounded sequence $(x = x_k)$ is said to be strongly σ -convergent to L if

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} |x_{\sigma^k(m)} - L| = 0 \text{ uniformly in } m$$

and in this case, we write $x_k \rightarrow L[V_\sigma]$. By $[V_\sigma]$, we denote the set of all strongly σ -convergent sequences.

In the case $\sigma(n) = n + 1$, the space $[V_\sigma]$ is the set of strongly almost convergent sequences $[\hat{C}]$.

Definitions and Notations

The concept of strong σ -convergence was generalized by Savaş [26] as below:

$$[V_\sigma]_p = \left\{ x = (x_k) : \lim_{m \rightarrow \infty} \frac{1}{m} \sum_{k=1}^m |x_{\sigma^k(n)} - L|^p = 0 \text{ uniformly in } n \right\},$$

where $0 < p < \infty$. If $p = 1$, then $[V_\sigma]_p = [V_\sigma]$. It is known that $[V_\sigma]_p \subset l_\infty$.

Definitions and Notations

A sequence $x = (x_k)$ is σ -statistically convergent to L if for every $\varepsilon > 0$,

$$\lim_{m \rightarrow \infty} \frac{1}{m} \left| \{k \leq m : |x_{\sigma^k(n)} - L| \geq \varepsilon\} \right| = 0,$$

uniformly in n . In this case we write $S_\sigma - \lim x = L$ or $x_k \rightarrow L(S_\sigma)$.

Definitions and Notations

Let $A \subseteq \mathbb{N}$ and

$$s_n := \min_m |A \cap \{\sigma(m), \sigma^2(m), \dots, \sigma^n(m)\}|$$

and

$$S_n := \max_m |A \cap \{\sigma(m), \sigma^2(m), \dots, \sigma^n(m)\}|.$$

If the following limits exist

$$\underline{V}(A) := \lim_{n \rightarrow \infty} \frac{s_n}{n}, \quad \overline{V}(A) := \lim_{n \rightarrow \infty} \frac{S_n}{n}$$

then they are called a lower and an upper σ -uniform density of the set A , respectively. If $\underline{V}(A) = \overline{V}(A)$, then $V(A) = \underline{V}(A) = \overline{V}(A)$ is called the σ -uniform density of A .

Definitions and Notations

Denote by \mathcal{I}_σ the class of all $A \subseteq \mathbb{N}$ with $V(A) = 0$.

A sequence (x_k) is said to be \mathcal{I}_σ -convergent to the number L if for every $\varepsilon > 0$ $A_\varepsilon = \{k : |x_k - L| \geq \varepsilon\} \in \mathcal{I}_\sigma$, that is, $V(A_\varepsilon) = 0$. In this case, we write $\mathcal{I}_\sigma - \lim x_k = L$.

Let (X, ρ) be a separable metric space. For any point $x \in X$ and any non-empty subset A of X , we define the distance from x to A by

$$d(x, A) = \inf_{a \in A} \rho(x, a).$$

Definitions and Notations

Throughout the paper, we let (X, ρ) be a separable metric space and A, A_{kj} be any non-empty closed subsets of X .

A double sequence $\{A_{kj}\}$ is Wijsman convergent to A if

$$P - \lim_{k,j \rightarrow \infty} d(x, A_{kj}) = d(x, A) \quad \text{or} \quad \lim_{k,j \rightarrow \infty} d(x, A_{kj}) = d(x, A)$$

for each $x \in X$. In this case, we write $W_2 - \lim A_{kj} = A$.

A double sequence of sets $\{A_{kj}\}$ is \mathcal{I}_{W_2} -convergent to A if for each $x \in X$ and for every $\varepsilon > 0$, $\{(k, j) \in \mathbb{N} \times \mathbb{N} : |d(x, A_{kj}) - d(x, A)| \geq \varepsilon\} \in \mathcal{I}_2$. In this case, we write $\mathcal{I}_{W_2} - \lim_{k,j \rightarrow \infty} d(x, A_{kj}) = d(x, A)$.

Definitions and Notations

A double sequence of sets $\{A_{kj}\}$ is $\mathcal{I}_{W_2}^*$ -convergent to A if there exists a set $M_2 \in \mathcal{F}(\mathcal{I}_2)$ (i.e., $\mathbb{N} \times \mathbb{N} \setminus M_2 = H \in \mathcal{I}_2$) such that for each $x \in X$

$$\lim_{\substack{k,j \rightarrow \infty \\ (k,j) \in M_2}} d(x, A_{kj}) = d(x, A).$$

In this case, we write $\mathcal{I}_{W_2}^* - \lim_{k,j \rightarrow \infty} d(x, A_{kj}) = d(x, A)$.

A double sequence of sets $\{A_{kj}\}$ is \mathcal{I}_2 -Cauchy sequence if for each $x \in X$ and for every $\varepsilon > 0$, there exists (p, q) in $\mathbb{N} \times \mathbb{N}$ such that $\{(k, j) \in \mathbb{N} \times \mathbb{N} : |d(x, A_{kj}) - d(x, A_{pq})| \geq \varepsilon\} \in \mathcal{I}_2$.

Definitions and Notations

A double sequence of sets $\{A_{kj}\}$ is $\mathcal{I}_{W_2}^*$ -Cauchy if there exists a set $M_2 \in \mathcal{F}(\mathcal{I}_2)$ ($\mathbb{N} \times \mathbb{N} \setminus M_2 = H \in \mathcal{I}_2$) such that for each $x \in X$,

$$\lim_{k,j,p,q \rightarrow \infty} |d(x, A_{kj}) - d(x, A_{pq})| = 0, \text{ for } (k, j), (p, q) \in M_2.$$

A double sequence $\{A_{kj}\}$ is said to be bounded if $\sup_{k,j} d(x, A_{kj}) < \infty$, for each $x \in X$. The set of all bounded double sequences of sets will be denoted by L_{∞}^2 .

Definitions and Notations

A sequence $\{A_k\}$ is said to be Wijsman invariant convergent to A if for each $x \in X$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n d(x, A_{\sigma^k(m)}) = d(x, A), \text{ uniformly in } m.$$

A sequence $\{A_k\}$ is said to be Wijsman strongly invariant convergent to A if for each $x \in X$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n |d(x, A_{\sigma^k(m)}) - d(x, A)| = 0, \text{ uniformly in } m.$$

A sequence $\{A_k\}$ is said to be Wijsman invariant statistical convergent to A if for every $\varepsilon > 0$ and for each $x \in X$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} |\{0 \leq k \leq n : |d(x, A_{\sigma^k(m)}) - d(x, A)| \geq \varepsilon\}| = 0,$$

uniformly in m .

Definitions and Notations

A sequence $\{A_k\}$ is said to be Wijsman p -strongly invariant convergent to A if for each $x \in X$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n |d(x, A_{\sigma^k(m)}) - d(x, A)|^p = 0, \text{ uniformly in } m,$$

where $0 < p < \infty$.

A sequence $\{A_k\}$ is said to be Wijsman \mathcal{I} -invariant convergent or \mathcal{I}_σ^W -convergent to A if for every $\varepsilon > 0$,

$A_\varepsilon = \{k : |d(x, A_k) - d(x, A)| \geq \varepsilon\} \in \mathcal{I}_\sigma$ that is, $V(A_\varepsilon) = 0$. In this case, we write $A_k \rightarrow A(\mathcal{I}_\sigma^W)$ and the set of all Wijsman \mathcal{I} -invariant convergent sequences of sets will be denoted \mathcal{I}_σ^W .

Definitions and Notations

An admissible ideal $\mathcal{I}_2 \subset 2^{\mathbb{N} \times \mathbb{N}}$ satisfies the property (AP2) if for every countable family of mutually disjoint sets $\{E_1, E_2, \dots\}$ belonging to \mathcal{I}_2 , there exists a countable family of sets $\{F_1, F_2, \dots\}$ such that $E_j \Delta F_j \in \mathcal{I}_2^0$, i.e., $E_j \Delta F_j$ is included in the finite union of rows and columns in $\mathbb{N} \times \mathbb{N}$ for each $j \in \mathbb{N}$ and $F = \bigcup_{j=1}^{\infty} F_j \in \mathcal{I}_2$ (hence $F_j \in \mathcal{I}_2$ for each $j \in \mathbb{N}$).

Main Result

In this section, we study the concepts of Wijsman invariant convergence, Wijsman invariant statistical convergence, Wijsman \mathcal{I}_2 -invariant convergence, Wijsman \mathcal{I}_2^* -invariant convergence, Wijsman p -strongly invariant convergence double sequence of sets and investigate the relationships among Wijsman invariant convergence, $[W_2 V_\sigma]_p$, $\mathcal{I}_{W_2}^\sigma$ and $\mathcal{I}_{W_2}^{*\sigma}$. Also, we introduce the concepts of $\mathcal{I}_{W_2}^\sigma$ -Cauchy double sequence and $\mathcal{I}_{W_2}^{*\sigma}$ -Cauchy double sequence of sets.

Main Result

Definition 1

Let $A \subseteq \mathbb{N} \times \mathbb{N}$ and

$$s_{mn} := \min_{k,j} |A \cap \{(\sigma(k), \sigma(j)), (\sigma^2(k), \sigma^2(j)), \dots, (\sigma^m(k), \sigma^n(j))\}|$$

and

$$S_{mn} := \max_{k,j} |A \cap \{(\sigma(k), \sigma(j)), (\sigma^2(k), \sigma^2(j)), \dots, (\sigma^m(k), \sigma^n(j))\}|.$$

If the following limits exists

$$\underline{V}_2(A) := \lim_{m,n \rightarrow \infty} \frac{s_{mn}}{mn}, \quad \overline{V}_2(A) := \lim_{m,n \rightarrow \infty} \frac{S_{mn}}{mn}$$

then they are called a lower and an upper σ -uniform density of the set A , respectively. If $\underline{V}_2(A) = \overline{V}_2(A)$, then $V_2(A) = \underline{V}_2(A) = \overline{V}_2(A)$ is called the σ -uniform density of A .

Main Result

Definition 2

A double sequence $\{A_{kj}\}$ is said to be Wijsman invariant convergent to A if for each $x \in X$,

$$\lim_{m,n \rightarrow \infty} \frac{1}{mn} \sum_{k,j=1,1}^{m,n} d(x, A_{\sigma^k(s), \sigma^j(t)}) = d(x, A), \text{ uniformly in } s, t.$$

Main Result

Definition 3

A double sequence $\{A_{kj}\}$ is said to be Wijsman \mathcal{I}_2 -invariant convergent or $\mathcal{I}_{W_2}^\sigma$ -convergent to A , if for every $\varepsilon > 0$,

$$A(\varepsilon, x) = \{(k, j) : |d(x, A_{kj}) - d(x, A)| \geq \varepsilon\} \in \mathcal{I}_2^\sigma$$

that is, $V_2(A(\varepsilon, x)) = 0$. In this case, we write $A_{kj} \rightarrow A(\mathcal{I}_{W_2}^\sigma)$ and the set of all Wijsman \mathcal{I}_2 -invariant convergent double sequences of sets will be denoted by $\mathcal{I}_{W_2}^\sigma$.

Main Result

Definition 4

Let $\mathcal{I}_2^\sigma \subset 2^{\mathbb{N} \times \mathbb{N}}$ be a strongly admissible ideal. A double sequence $\{A_{kj}\}$ is Wijsman \mathcal{I}_2^* -invariant convergent or $\mathcal{I}_{W_2}^{*\sigma}$ -convergent to A if and only if there exists a set $M_2 \in \mathcal{F}(\mathcal{I}_2^\sigma)$ ($\mathbb{N} \times \mathbb{N} \setminus M_2 = H \in \mathcal{I}_2^\sigma$) such that for each $x \in X$,

$$\lim_{\substack{k,j \rightarrow \infty \\ (k,j) \in M_2}} d(x, A_{kj}) = d(x, A).$$

Main Result

Theorem 5

Let $\mathcal{I}_2^\sigma \subset 2^{\mathbb{N} \times \mathbb{N}}$ be a strongly admissible ideal. If a sequence $\{A_{kj}\}$ is $\mathcal{I}_{W_2}^{\sigma}$ -convergent to A , then this sequence is $\mathcal{I}_{W_2}^\sigma$ -convergent to A .*

Main Result

Proof: Since $\mathcal{I}_{W_2}^{*\sigma} - \lim_{k,j \rightarrow \infty} d(x, A_{kj}) = d(x, A)$, there exists a set $M_2 \in \mathcal{F}(\mathcal{I}_2^\sigma)$ ($\mathbb{N} \times \mathbb{N} \setminus M_2 = H \in \mathcal{I}_2^\sigma$) such that for each $x \in X$,

$$\lim_{\substack{k,j \rightarrow \infty \\ (k,j) \in M_2}} d(x, A_{kj}) = d(x, A).$$

Main Result

Let $\varepsilon > 0$. Then, there exists $k_0 \in \mathbb{N}$ such that for each $x \in X$,

$$|d(x, A_{kj}) - d(x, A)| < \varepsilon,$$

for all $(k, j) \in M_2$ and $k, j \geq k_0$. Hence, for every $\varepsilon > 0$ and each $x \in X$, we have

$$\begin{aligned} T(\varepsilon, x) &= \{(k, j) \in \mathbb{N} \times \mathbb{N} : |d(x, A_{kj}) - d(x, A)| \geq \varepsilon\} \\ &\subset H \cup \left(M_2 \cap \left((\{1, 2, \dots, (k_0 - 1)\} \times \mathbb{N}) \cup (\mathbb{N} \times \{1, 2, \dots, (k_0 - 1)\}) \right) \right). \end{aligned}$$

Main Result

Since $\mathcal{I}_2^\sigma \subset 2^{\mathbb{N} \times \mathbb{N}}$ is a strongly admissible ideal,

$$H \cup \left(M_2 \cap \left((\{1, 2, \dots, (k_0 - 1)\} \times \mathbb{N}) \cup (\mathbb{N} \times \{1, 2, \dots, (k_0 - 1)\}) \right) \right) \in \mathcal{I}_2^\sigma,$$

so we have $T(\varepsilon, x) \in \mathcal{I}_2^\sigma$ that is $V_2(T(\varepsilon, x)) = 0$. Hence,

$$\mathcal{I}_{W_2}^\sigma - \lim_{k, j \rightarrow \infty} d(x, A_{kj}) = d(x, A).$$

Main Result

Theorem 6

Let $\mathcal{I}_2^\sigma \subset 2^{\mathbb{N} \times \mathbb{N}}$ be a strongly admissible ideal with property (AP2). If $\{A_{kj}\}$ is $\mathcal{I}_{W_2}^\sigma$ -convergent to A , then $\{A_{kj}\}$ is $\mathcal{I}_{W_2}^{*\sigma}$ -convergent to A .

Main Result

Proof: Suppose that \mathcal{I}_2^σ satisfies property (AP2). Let $\{A_{kj}\}$ is $\mathcal{I}_{W_2}^\sigma$ -convergent to A . Then,

$$T(\varepsilon, x) = T_\varepsilon = \{(k, j) \in \mathbb{N} \times \mathbb{N} : |d(x, A_{kj}) - d(x, A)| \geq \varepsilon\} \in \mathcal{I}_2^\sigma \quad (4.1)$$

for every $\varepsilon > 0$ and for each $x \in X$. Put

$$T_1 = T(1, x) = \{(k, j) \in \mathbb{N} \times \mathbb{N} : |d(x, A_{kj}) - d(x, A)| \geq 1\}$$

and

$$T_\nu = T(\nu, x) = \left\{ (k, j) \in \mathbb{N} \times \mathbb{N} : \frac{1}{\nu} \leq |d(x, A_{kj}) - d(x, A)| < \frac{1}{\nu - 1} \right\}$$

for $\nu \geq 2$ and $\nu \in \mathbb{N}$.

Main Result

Obviously $T_i \cap T_j = \emptyset$ for $i \neq j$ and $T_i \in \mathcal{I}_2^\sigma$ for each $i \in \mathbb{N}$. By property (AP2) there exists a sequence of sets $\{E_\nu\}_{\nu \in \mathbb{N}}$ such that $T_i \Delta E_i$ is included in finite union of rows and columns in $\mathbb{N} \times \mathbb{N}$ for each i and

$$E = \bigcup_{i=1}^{\infty} E_i \in \mathcal{I}_2^\sigma.$$

We shall prove that for $M_2 = \mathbb{N} \times \mathbb{N} \setminus E$ we have

$$\lim_{\substack{k,j \rightarrow \infty \\ (k,j) \in M_2}} d(x, A_{kj}) = d(x, A).$$

Main Result

Let $\eta > 0$ be given. Choose $\nu \in \mathbb{N}$ such that $\frac{1}{\nu} < \eta$. Then,

$$\{(k, j) \in \mathbb{N} \times \mathbb{N} : |d(x, A_{kj}) - d(x, A)| \geq \eta\} \subset \bigcup_{i=1}^{\nu} T_i$$

Main Result

Since $T_i \Delta E_i$, $i = 1, 2, \dots$ are included in finite union of rows and columns, there exists $n_0 \in \mathbb{N}$ such that

$$\left(\bigcup_{i=1}^{\vee} T_i \right) \cap \{(k, j) : k \geq n_0 \wedge j \geq n_0\} = \left(\bigcup_{i=1}^{\vee} E_i \right) \cap \{(k, j) : k \geq n_0 \wedge j \geq n_0\}. \quad (4.2)$$

If $k, j > n_0$ and $(k, j) \notin E$, then

$$(k, j) \notin \bigcup_{i=1}^{\vee} E_i \text{ and } (k, j) \notin \bigcup_{i=1}^{\vee} T_i.$$

This implies that

$$|d(x, A_{kj}) - d(x, A)| < \frac{1}{\vee} < \eta.$$

Hence, we have

$$\lim_{\substack{k, j \rightarrow \infty \\ (k, j) \in M_2}} d(x, A_{kj}) = d(x, A).$$

Main Result

Definition 7

A double sequence $\{A_{kj}\}$ is said to be Wijsman \mathcal{I}_2 -invariant Cauchy sequence or $\mathcal{I}_{W_2}^\sigma$ -Cauchy sequence, if for every $\varepsilon > 0$ and for each $x \in X$, there exist numbers $r = r(\varepsilon, x), s = s(\varepsilon, x) \in \mathbb{N}$ such that

$$A(\varepsilon, x) = \{(k, j) : |d(x, A_{kj}) - d(x, A_{rs})| \geq \varepsilon\} \in \mathcal{I}_2^\sigma,$$

that is, $V_2(A(\varepsilon, x)) = 0$.

Main Result

Definition 8

A double sequence $\{A_{kj}\}$ is $\mathcal{I}_{W_2}^{*\sigma}$ -Cauchy if there exists a set $M_2 \in \mathcal{F}(\mathcal{I}_2^\sigma)$ (i.e., $\mathbb{N} \times \mathbb{N} \setminus M_2 = H \in \mathcal{I}_2^\sigma$) such that for every $x \in X$ and $(k, j), (p, q) \in M_2$

$$\lim_{k,j,p,q \rightarrow \infty} |d(x, A_{kj}) - d(x, A_{pq})| = 0.$$

We give following theorems which show relationships between $\mathcal{I}_{W_2}^\sigma$ -convergence, $\mathcal{I}_{W_2}^\sigma$ -Cauchy sequence and $\mathcal{I}_{W_2}^{*\sigma}$ -Cauchy sequence. The proof of them are similar to the proof of Theorems in [5, 18], so we omit them.

Main Result

Theorem 9

If a double sequence $\{A_{kj}\}$ is $\mathcal{I}_{W_2}^\sigma$ -convergent, then $\{A_{kj}\}$ is an $\mathcal{I}_{W_2}^\sigma$ -Cauchy double sequence of sets.

Theorem 10

If a double sequence $\{A_{kj}\}$ is $\mathcal{I}_{W_2}^{\sigma}$ -Cauchy double sequence, then $\{A_{kj}\}$ is $\mathcal{I}_{W_2}^\sigma$ -Cauchy double sequence of sets.*

Theorem 11

Let \mathcal{I}_2^σ has property (AP2). Then, the concepts $\mathcal{I}_{W_2}^\sigma$ -Cauchy sequence and $\mathcal{I}_{W_2}^{\sigma}$ -Cauchy sequence of sets coincides.*

Main Result

Definition 12

A double sequence $\{A_{kj}\}$ is said to be Wijsman strongly invariant convergent to A , if for each $x \in X$,

$$\lim_{m,n \rightarrow \infty} \frac{1}{mn} \sum_{k,j=1,1}^{m,n} |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)| = 0, \text{ uniformly in } s, t.$$

Main Result

Definition 13

A double sequence $\{A_{kj}\}$ is said to be Wijsman p -strongly invariant convergent to A , if for each $x \in X$,

$$\lim_{m,n \rightarrow \infty} \frac{1}{mn} \sum_{k,j=1,1}^{m,n} |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)|^p = 0, \text{ uniformly in } s, t.$$

where $0 < p < \infty$. In this case, we write $A_k \rightarrow A[W_2V_\sigma]_p$ and the set of all Wijsman p -strongly invariant convergent sequences of sets will be denoted by $[W_2V_\sigma]_p$.

Main Result

Theorem 14

Let $\{A_{kj}\}$ is bounded sequence. If $\{A_{kj}\}$ is $\mathcal{I}_{W_2}^\sigma$ -convergent to A , then $\{A_{kj}\}$ is Wijsman invariant convergent to A .

Main Result

Proof: Let $m, n \in \mathbb{N}$ be arbitrary and $\varepsilon > 0$. For each $x \in X$, we estimate

$$u(s, t, m, n, x) = \left| \frac{1}{mn} \sum_{k,j=1,1}^{m,n} d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A) \right|.$$

Then, for each $x \in X$ we have

$$u(s, t, m, n, x) \leq u^1(s, t, m, n, x) + u^2(s, t, m, n, x)$$

Main Result

where

$$u^1(s, t, m, n, x) = \frac{1}{mn} \sum_{\substack{k,j=1,1 \\ |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)| \geq \varepsilon}}^{m,n} |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)|$$

and

$$u^2(s, t, m, n, x) = \frac{1}{mn} \sum_{\substack{k,j=1,1 \\ |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)| < \varepsilon}}^{m,n} |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)|.$$

Main Result

Therefore, we have

$$u^2(s, t, m, n, x) < \varepsilon,$$

for each $x \in X$ and for every $s, t = 1, 2, \dots$. The boundedness of $\{A_{kj}\}$ implies that there exists $L > 0$ such that

$$|d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)| \leq L, \quad (k, s, j, t = 1, 2, \dots),$$

then this implies that

$$\begin{aligned} u^1(s, t, m, n, x) &\leq \frac{L}{mn} |\{1 \leq k \leq m, 1 \leq j \leq n : |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)| \geq \varepsilon\}| \\ &\leq L \frac{\max_{s,t} |\{1 \leq k \leq m, 1 \leq j \leq n : |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)| \geq \varepsilon\}|}{mn} \\ &= L \frac{S_{mn}}{mn}. \end{aligned}$$

Hence $\{A_{kj}\}$ is Wijsman invariant convergent to A .

Main Result

Theorem 15

Let $\mathcal{I}_2^\sigma \subset 2^{\mathbb{N} \times \mathbb{N}}$ be a strongly admissible ideal and $0 < p < \infty$.

- (i) If $A_{kj} \rightarrow A([W_2 V_\sigma]_p)$, then $A_{kj} \rightarrow A(\mathcal{I}_{W_2}^\sigma)$.
- (ii) If $\{A_{kj}\} \in L_\infty^2$ and $A_{kj} \rightarrow A(\mathcal{I}_{W_2}^\sigma)$, then $A_{kj} \rightarrow A([W_2 V_\sigma]_p)$.
- (iii) If $\{A_{kj}\} \in L_\infty^2$, then $\{A_{kj}\}$ is $\mathcal{I}_{W_2}^\sigma$ -convergent to A if and only if $A_{kj} \rightarrow A([W_2 V_\sigma]_p)$.

Main Result

(i) : Assume that $A_{kj} \rightarrow A([W_2 V_\sigma]_p)$, for every $\varepsilon > 0$ and for each $x \in X$. Then, we can write

$$\begin{aligned}
 & \sum_{k,j=1,1}^{m,n} |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)|^p \\
 & \geq \sum_{k,j=1,1}^{m,n} |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)|^p \\
 & \quad |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)| \geq \varepsilon \\
 & \geq \varepsilon^p |\{k \leq m, j \leq n : |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)| \geq \varepsilon\}| \\
 & \geq \varepsilon^p \max_{s,t} |\{k \leq m, j \leq n : |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)| \geq \varepsilon\}|
 \end{aligned}$$

Main Result

and

$$\begin{aligned}
 & \frac{1}{mn} \sum_{k,j=1,1}^{m,n} |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)|^p \\
 \geq & \varepsilon^p \cdot \frac{\max_{s,t} |\{k \leq m, j \leq n : |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)| \geq \varepsilon\}|}{mn} \\
 = & \varepsilon^p \frac{S_{mn}}{mn}
 \end{aligned}$$

Main Result

for every $s, t = 1, 2, \dots$. This implies

$$\lim_{m, n \rightarrow \infty} \frac{S_{mn}}{mn} = 0$$

and so $\{A_{kj}\}$ is $(\mathcal{I}_{W_2}^\sigma)$ -convergent to A .

Main Result

(ii): Suppose that $\{A_{kj}\} \in L_{\infty}^2$ and $A_{kj} \rightarrow A(\mathcal{I}_{W_2}^{\sigma})$. Let $0 < p < \infty$ and $\varepsilon > 0$. By assumption we have $V_2(A_{\varepsilon}) = 0$. Since $\{A_{kj}\}$ is bounded, $\{A_{kj}\}$ implies that there exists $L > 0$ such that for each $x \in X$,

$$|d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)| \leq L$$

for all k, s, j and t .

Main Result

Then, we have

$$\begin{aligned}
 & \frac{1}{mn} \sum_{k,j=1,1}^{m,n} |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)|^p \\
 = & \frac{1}{mn} \sum_{\substack{k,j=1,1 \\ |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)| \geq \varepsilon}}^{m,n} |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)|^p \\
 + & \frac{1}{mn} \sum_{\substack{k,j=1,1 \\ |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)| < \varepsilon}}^{m,n} |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)|^p \\
 \leq & L \frac{\max_{s,t} |\{k \leq m, j \leq n : |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)| \geq \varepsilon\}|}{mn} + \varepsilon^p \\
 \leq & L \frac{S_{mn}}{mn} + \varepsilon^p,
 \end{aligned}$$

Main Result

for each $x \in X$ we obtain

$$\lim_{m,n \rightarrow \infty} \frac{1}{mn} \sum_{k,j=1,1}^{m,n} |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)|^p = 0, \text{ uniformly in } s, t.$$

(iii) : This is immediate consequence of (i) and (ii).

Main Result

Definition 16

A double sequence $\{A_{kj}\}$ is said to be Wijsman invariant statistical convergent or W_2S_σ -convergent to A , if for every $\varepsilon > 0$ and for each $x \in X$,

$$\lim_{m,n \rightarrow \infty} \frac{1}{mn} |\{k \leq m, j \leq n : |d(x, A_{\sigma^k(s), \sigma^j(t)}) - d(x, A)| \geq \varepsilon\}| = 0,$$

uniformly in s,t.

Theorem 17

A sequence $\{A_{kj}\}$ is W_2S_σ -convergent to A if and only if it is $\mathcal{I}_{W_2}^\sigma$ -convergent to A .

References I

- [1] M. Baronti, and P. Papini, *Convergence of sequences of sets*, In: Methods of functional analysis in approximation theory, ISNM 76, Birkhauser-Verlag, Basel, pp. 133-155, (1986).
- [2] G. Beer, *On convergence of closed sets in a metric space and distance functions*, Bull. Aust. Math. Soc. **31** (1985), 421–432.
- [3] G. Beer, *Wijsman convergence: A survey*, Set-Valued Var. Anal. **2** (1994), 77–94.
- [4] P. Das, P. Kostyrko, W. Wilczyński, P. Malik, *I and I^* -convergence of double sequences*, Math. Slovaca, **58** (5) (2008), 605–620.
- [5] E. Dndar, B. Altay, *\mathcal{I}_2 -convergence and \mathcal{I}_2 -Cauchy of double sequences*, Acta Mathematica Scientia, **34B**(2) (2014), 343–353.
- [6] H. Fast, *Sur la convergence statistique*, Colloq. Math. **2** (1951) 241–244.

References II

- [7] Kişi, Ö. and Nuray, F. *A new convergence for sequences of sets*, Abstract and Applied Analysis, vol. 2013, Article ID 852796, 6 pages. <http://dx.doi.org/10.1155/2013/852796>.
- [8] P. Kostyrko, T. Šalát, W. Wilczyński, *\mathcal{I} -Convergence*, Real Anal. Exchange **26**(2) (2000) 669–686.
- [9] V. Kumar, *On I and I^* -convergence of double sequences*, Math. Commun. **12** (2007), 171–181.
- [10] G. Lorentz, *A contribution to the theory of divergent sequences*, Acta Math. **80** (1948) 167–190.
- [11] M. Mursaleen, O. H. H. Edely, *On the invariant mean and statistical convergence*, Appl. Math. Lett. **22** (2009) 1700–1704.
- [12] Mursaleen, O. H. H. Edely, *Statistical convergence of double sequences*, J. Math. Anal. Appl. **288** (2003) 223–231.

References III

- [13] M. Mursaleen, *Matrix transformation between some new sequence spaces*, Houston J. Math. **9** (1983) 505–509.
- [14] M. Mursaleen, *On finite matrices and invariant means*, Indian J. Pure and Appl. Math. **10** (1979) 457–460.
- [15] A. Nabiev, S. Pehlivan, M. Gürdal, *On \mathcal{I} -Cauchy sequences*, Taiwanese J. Math. **11**(2) (2007) 569–5764.
- [16] F. Nuray, B. E. Rhoades, *Statistical convergence of sequences of sets*, Fasc. Math. **49** (2012), 87–99.
- [17] F. Nuray, W.H. Ruckle, *Generalized statistical convergence and convergence free spaces*, J. Math. Anal. Appl. **245** (2000), 513–527.
- [18] F. Nuray, E. Dndar, U. Ulusu, *Wijsman \mathcal{I}_2 -convergence of double sequences of closed sets*, Pure and Applied Mathematics Letters, **2** (2014), 35-39.

References IV

- [19] F. Nuray, E. Savaş, *Invariant statistical convergence and A -invariant statistical convergence*, Indian J. Pure Appl. Math. **10** (1994) 267–274.
- [20] F. Nuray, H. Gök, U. Ulusu, *\mathcal{I}_σ -convergence*, Math. Commun. **16** (2011) 531–538.
- [21] U. Ulusu, F. Nuray, *Lacunary \mathcal{I}_σ -convergence*, (Under Communication).
- [22] N. Pancaroğlu, F. Nuray, *Statistical lacunary invariant summability*, Theoretical Mathematics and Applications **3**(2) (2013) 71–78.
- [23] N. Pancaroğlu, F. Nuray, *On Invariant Statistically Convergence and Lacunary Invariant Statistically Convergence of Sequences of Sets*, Progress in Applied Mathematics, **5**(2) (2013), 23–29.

References V

- [24] N. Pancaroğlu, E. Dndar and F. Nuray, *Wijsman \mathcal{I} -Invariant Convergence of Sequences of Sets*, (Under Communication).
- [25] R. A. Raimi, *Invariant means and invariant matrix methods of summability*, Duke Math. J. **30** (1963) 81–94.
- [26] E. Savaş, *Some sequence spaces involving invariant means*, Indian J. Math. **31** (1989) 1–8.
- [27] E. Savaş, *Strong σ -convergent sequences*, Bull. Calcutta Math. **81** (1989) 295–300.
- [28] E. Savaş, F. Nuray, *On σ -statistically convergence and lacunary σ -statistically convergence*, Math. Slovaca **43**(3) (1993) 309–315.
- [29] P. Schaefer, *Infinite matrices and invariant means*, Proc. Amer. Math. Soc. **36** (1972) 104–110.

References VI

- [30] I. J. Schoenberg, *The integrability of certain functions and related summability methods*, Amer. Math. Monthly **66** (1959) 361–375.
- [31] Y. Sever, U. Ulusu and E. Dndar, *On Strongly I and I^* -Lacunary Convergence of Sequences of Sets*, AIP Conference Proceedings, 1611, 357 (2014); doi: 10.1063/1.4893860, 7 pages.
- [32] Y. Sever, . Talo, *On Statistical Convergence of Double Sequences of Closed Sets*, Filomat **30**(3) (2016), 533–539, DOI 10.2298/FIL1603533S.
- [33] Y. Sever, . Talo, B. Altay, *On convergence of double sequences of closed sets*, Contemp. Anal. Appl. Math. **3** (2015), 30–49.
- [34] . Talo, Y. Sever, F. Başar, *On statistically convergent sequences of closed set*, Filomat, **30**(6) (2016), (In press).

References VII

- [35] U. Ulusu, F. Nuray, *Lacunary statistical convergence of sequence of sets*, Progress in Applied Mathematics, **4**(2) (2012), 99–109.
- [36] U. Ulusu and E. Dndar, *\mathcal{I} -Lacunary Statistical Convergence of Sequences of Sets*, Filomat, **28**(8) (2013), 1567–1574.
- [37] U. Ulusu, F. Nuray, *On Strongly Lacunary Summability of Sequences of Sets*, Journal of Applied Mathematics and Bioinformatics, **3**(3) (2013), 75–88.
- [38] R. A. Wijsman, *Convergence of sequences of convex sets, cones and functions*, Bull. Amer. Math. Soc. **70** (1964), 186–188.
- [39] R. A. Wijsman, *Convergence of Sequences of Convex sets, Cones and Functions II*, Trans. Amer. Math. Soc. **123** (1) (1966), 32–45.

Thanks for your attention.